

Caretta caretta (L.)—continued.

A.M.N.H. No.	(a) Length Carapace	(b) Breadth	(c) Length Plastron	b a	c a
4610	47	38	36	.81	.76
4611	47	39	35	.83	.74
4612	47	37	35	.79	.74
4613	48	37	37	.77	.77

KARL P. SCHMIDT,
EMMETT R. DUNN,
New York, N. Y.

FURTHER NOTES ON PSEUDEMYS AT PLYMOUTH, MASSACHUSETTS.

The two short articles in the December "COPEIA" having stimulated some new interest in this connection, I have to record the following new ponds in Plymouth County, Massachusetts, in which *Pseudemys rubriventris* has been observed: Great South Pond and Crooked Pond. The Island Pond mentioned in my notes is the one about one-half mile northeast of Gunners Exchange. Hallfield Pond, the existence of which Dr. Lucas doubted, is a small four-acre pond between Hoyt and Boot Ponds. *P. rubriventris* does not occur in Half-way Pond as far as I know. It is a curious fact that in the ponds where this turtle is found, it seems to be the most common species. It is also of interest that the "Plymouth Sabbatia" and other plants, as well as certain insects, notably dragon flies, which are distributed over the same local areas as this species of turtle, are all southern forms of life.

HAROLD L. BABCOCK,
Boston, Mass.